

K^0 - $\bar{K}^0$ mixing from the Schrödinger functional and twisted mass QCD*

M. Guagnelli^a, J. Heitger^b, C. Pena^a, S. Sint^c and A. Vladikas^a

^aINFN Sezione Roma II, c/o Dipartimento di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy.

^bWestfälische Wilhelms-Universität Münster, Institut für Theoretische Physik,
Wilhelm-Klemm-Straße 9, D-48149 Münster, Germany.

^cCERN, Theory Division, CH-1211 Geneva 23, Switzerland.

We describe how the Schrödinger functional and twisted mass QCD can be used to compute the kaon B_K parameter on the lattice with Wilson fermions. This new approach is expected to reduce the systematic uncertainties on B_K through a better control of its (nonperturbative) renormalisation. Preliminary results for the bare matrix element in a physical volume of $1.5^3 \times 4.5 \text{ fm}^4$ are presented. The renormalisation of the matrix element in a Schrödinger functional scheme is also discussed.

1. INTRODUCTION

Out of the physical quantities related to CP violation in the Standard Model, the parameter ε accounting for indirect CP violation in the kaon system is the one with the simplest non-perturbative contribution. It is given by the kaon B -parameter B_K , defined as:

$$B_K = \frac{\langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle}{\frac{8}{3} F_K^2 m_K^2}, \quad (1)$$

where $O^{\Delta S=2}$ is the $\Delta S = 2$ operator of the effective weak Hamiltonian:

$$O^{\Delta S=2} = (\bar{s} \gamma_\mu^L d) (\bar{s} \gamma_\mu^L d), \quad (2)$$

and $\gamma_\mu^L = \frac{1}{2} \gamma_\mu (\mathbf{1} - \gamma_5)$. In an obvious notation, it is convenient to split $O^{\Delta S=2}$ in parity-even and parity-odd parts:

$$O^{\Delta S=2} = O_{VV+AA}^{\Delta S=2} - O_{VA+AV}^{\Delta S=2}. \quad (3)$$

Since parity is a QCD symmetry, only the parity-even part $O_{VV+AA}^{\Delta S=2}$ will yield a nonvanishing K^0 - $\bar{K}^0$ matrix element.

In spite of the absence of both power-divergent subtractions and final state interactions, the lattice computation of B_K is still a very difficult

problem. Besides the systematic error arising from the quenched approximation, the breaking of chirality (due to the use of Wilson fermions) leads to a mixing under renormalisation of $O^{\Delta S=2}$ with four other operators of dimension six. It is now understood that the corresponding mixing coefficients, as well as the overall, scale-dependent, renormalisation constant, require an accurate nonperturbative computation. This has been dealt with by using Ward identities and RI-MOM renormalisation [1,2], but the result still reflects large numerical uncertainties. On the other hand, the use of lattice actions with better chiral properties has yielded results which have not manifested clear agreement with those obtained with Wilson fermions. We refer the reader to [3] for reviews on the subject and references.

To improve on this situation, we propose a new computation of B_K with Wilson fermions. It adopts the Schrödinger functional (SF) [4] for the computation of scale-dependent renormalisation constants, and a twisted mass QCD (tmQCD) action [5], in order to avoid the mixing of $O^{\Delta S=2}$ with other operators under renormalisation. An alternative proposal for the elimination of this mixing has been presented in [6].

*Based on a talk by C. Pena at the International Symposium of Lattice Field Theory, August 19-24 2001, Berlin, Germany.

2. tmQCD FOR B_K

Twisted mass QCD was proposed as a means of avoiding the problem of exceptional configurations, allowing lattice QCD computations with light quark masses closer to the chiral limit [5]. For the computation of B_K we will use an action with a twisted doublet of light quarks $\psi = (u, d)$ and a standard (untwisted) strange quark s :

$$S = a^4 \sum_x [\bar{\psi}(x) (D + m_l + i\mu_l \gamma_5 \tau^3) \psi(x) + \bar{s}(x) (D + m_s) s(x)] , \quad (4)$$

where τ^3 acts on isospin indices and μ_l is the “twisted mass”. In the classical theory at the continuum limit, the standard QCD action is recovered by changing fermionic variables through an axial rotation of the form $\psi \rightarrow \exp\{-i\alpha\gamma_5\tau^3/2\}\psi$, where $\tan(\alpha) = \mu_l/m_l$. This rotation leaves invariant the combination $M_l = \sqrt{m_l^2 + \mu_l^2}$, which can therefore be regarded as the physical bare light quark mass; on the other hand, it induces a mapping between composite operators in the two theories. It has been shown in [5] that this classical equivalence carries over to appropriately renormalised correlation functions in the quantum theory. The twisting angle is then given by $\tan(\alpha) = \mu_{l,R}/m_{l,R}$, where the masses are renormalised.

By fixing renormalised masses so that $\alpha = \pi/2$, the mapping between operators induced by the corresponding axial rotation leads to the relation:

$$\begin{aligned} \langle \bar{K}^0 | O_{VA+AV}^{\Delta S=2} | K^0 \rangle_{\text{tmQCD}}^{\alpha=\pi/2} &= \\ &= i \langle \bar{K}^0 | O_{VV+AA}^{\Delta S=2} | K^0 \rangle_{\text{QCD}} \\ &= i \langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle_{\text{QCD}} \end{aligned} \quad (5)$$

(the above is to be understood as valid for renormalised matrix elements). Parity violation at nonzero twisted mass μ_l ensures that the tmQCD matrix element does not vanish. Computing the lhs of Eq. (5), rather than the rhs, is advantageous, as the parity-odd operator $O_{VA+AV}^{\Delta S=2}$ is protected by $\mathcal{CP}\mathcal{S}$ symmetry from mixing with other operators, and hence it renormalises in a purely multiplicative fashion [7]. This is clearly true if renormalisation is carried out in the chiral limit,

where tmQCD and QCD (and their symmetries) coincide.

We show the feasibility of a direct computation of the matrix element of $O_{VA+AV}^{\Delta S=2}$ with the tmQCD action and SF boundary conditions by presenting preliminary results for the bare value of B_K , obtained on a $16^3 \times 48$ lattice at $\beta = 6.0$. This corresponds approximately to a physical volume of $1.5^3 \times 4.5 \text{ fm}^4$. In the computation we have kept the s and d masses degenerate and close to the physical s quark mass, as done in previous computations of B_K . We show in Fig. 1 the ratio of correlations:

$$R(x_0) = \frac{-i \langle \mathcal{O}'_{ds} O_{VA+AV}^{\Delta S=2}(x) \mathcal{O}_{ds} \rangle}{\frac{8}{3} \langle \mathcal{O}'_{ds} \mathcal{K}(x) \rangle \langle \mathcal{K}(x) \mathcal{O}_{ds} \rangle} , \quad (6)$$

where $\mathcal{K} = -\frac{i}{\sqrt{2}} \bar{s} \gamma_0 (1 + i\gamma_5) d$ is the operator obtained by axially rotating the QCD A_0 current with $\alpha = \pi/2$, and $\mathcal{O}, \mathcal{O}'$ are the standard SF pseudoscalar sources:

$$\mathcal{O}_{f_1 f_2} = L^{-3} \sum_{\mathbf{xy}} \bar{\zeta}_{f_1}(\mathbf{x}) \gamma_5 \zeta_{f_2}(\mathbf{y}) , \quad (7)$$

$$\mathcal{O}'_{f_1 f_2} = L^{-3} \sum_{\mathbf{xy}} \bar{\zeta}'_{f_1}(\mathbf{x}) \gamma_5 \zeta'_{f_2}(\mathbf{y}) , \quad (8)$$

where f_1, f_2 label two distinct flavours. The central plateau of R corresponds to the bare value of B_K . As expected, it appears in the time interval $0.35 \lesssim x_0/T \lesssim 0.65$, where the single kaon states coming from Euclidean times 0 and T are isolated (as shown by the effective mass plots of Fig. 1). Fitting the R -plateau to a constant we get $B_K(a^{-1} \sim 2 \text{ GeV}) = 0.96(2)$.

3. SF RENORMALISATION

The renormalisation constant of the matrix element at renormalisation scale $\mu = L^{-1}$ can be evaluated on symmetric lattices of physical volume L^4 . The SF renormalisation condition, which is imposed at the chiral limit $m_{l,R} = m_{s,R} = \mu_{l,R} = 0$, is then:

$$Z_{VA+AV}(g_0, L) \frac{F(L/2)}{f_1^{3/2}} = \frac{F(L/2)}{f_1^{3/2}} \bigg|_{g_0^2=0} , \quad (9)$$




Figure 1. Plateaux for B_K and effective kaon masses, from 250 configurations, for a $16^3 \times 48$ lattice at $\beta = 6.0$.

where f_1 is a wave function renormalisation factor, defined as:

$$f_1 = -\frac{1}{2} \langle \mathcal{O}'_{sd} \mathcal{O}_{ds} \rangle, \quad (10)$$

and F is a correlator involving $O_{VA+AV}^{\Delta S=2}$, chosen so as not to vanish in the chiral limit. We opt for the simple choice:

$$F(x_0) = L^6 \langle \mathcal{O}'_{us} O_{VA+AV}^{\Delta S=2}(x) \mathcal{O}_{ds} \mathcal{O}_{du} \rangle. \quad (11)$$

Once Z_{VA+AV} has been defined at given values of the bare coupling g_0 and of the renormalisation scale $\mu = L^{-1}$, a step scaling function that describes the (nonperturbative) running with μ can be defined at fixed value $u = \bar{g}^2(L)$ of the renormalised coupling as:

$$\sigma_{VA+AV}(u) = \lim_{a \rightarrow 0} \frac{Z_{VA+AV}(g_0, 2L)}{Z_{VA+AV}(g_0, L)} \Big|_{\bar{g}^2(L)=u}. \quad (12)$$

Having computed σ_{VA+AV} for a number of values of u , it is then possible to run the renormalised matrix element to large scales $\mu = L^{-1}$, at which the perturbative matching to other renormalisation schemes, such as $\overline{\text{MS}}$, can be performed.

By computing various Z constants, obtained from different sources $\mathcal{O}, \mathcal{O}'$ (i.e. different renormalisation conditions), we hope to get an idea of

the size of $O(a)$ effects, which arise due to the use of the unimproved four-quark operator.

4. CONCLUSIONS

We have described how the SF and tmQCD can be used to achieve a better control of the renormalisation properties of the kaon B -parameter B_K . A first feasibility study for the bare matrix element giving B_K in tmQCD has been presented, and the strategy for the SF renormalisation discussed.

We mention that the same techniques can be extended to all $\Delta S = 2$ operators of interest, thus avoiding the spurious mixing with operators of the same dimension induced by the use of Wilson fermions. The computation of the step scaling functions for the complete basis of such operators is already under way.

ACKNOWLEDGEMENTS

We wish to thank D. Bećirević, K. Jansen, G. Martinelli and R. Sommer for useful discussions. This work was supported in part by the European Community's Human Potential Programme under contract HPRN-CT-2000-00145, Hadrons/Lattice QCD. C.P. acknowledges the financial support provided through this contract.

REFERENCES

1. L. Conti et al., Phys. Lett. **B421** (1998) 273.
2. S. Aoki et al., Phys. Rev. **D60** (1999) 034511.
3. L. Lellouch, Nucl. Phys. B (Proc. Suppl.) **94** (2001) 142.
G. Martinelli, these proceedings.
4. M. Lüscher et al., Nucl. Phys. **B384** (1992) 168.
S. Sint, Nucl. Phys. **B421** (1994) 135.
5. R. Frezzotti et al., Nucl. Phys. B (Proc. Suppl.) **83** (2000) 941.
R. Frezzotti et al., JHEP **0108** (2001) 058.
6. D. Bećirević et al., Phys. Lett. **B487** (2000) 74; these proceedings, hep-lat/0110006.
7. C. Bernard et al., Nucl. Phys. B (Proc. Suppl.) **4** (1988) 483.
A. Donini et al., Eur. Phys. J. **C10** (1999) 121.